

Metric spaces and homotopy types

J.F. Jardine*

Department of Mathematics
University of Western Ontario
London, Ontario, Canada

jardine@uwo.ca

September 1, 2020

Abstract

By analogy with methods of Spivak, there is a realization functor which takes a persistence diagram Y in simplicial sets to an extended pseudo-metric space $Re(Y)$. The functor Re has a right adjoint, called the singular functor, which takes an ep-metric space Z to a persistence diagram $S(Z)$. We give an explicit description of $Re(Y)$, and show that it depends only on the 1-skeleton $sk_1 Y$ of Y . If X is a totally ordered ep-metric space, then there is an isomorphism $Re(V_*(X)) \cong X$, between the realization of the Vietoris-Rips diagram $V_*(X)$ and the ep-metric space X . The persistence diagrams $V_*(X)$ and $S(X)$ are sectionwise equivalent for all such X .

Introduction

An extended pseudo-metric space, here called an ep-metric space, is a set X together with a function $d : X \times X \rightarrow [0, \infty]$ such that the following conditions hold:

- 1) $d(x, x) = 0$,
- 2) $d(x, y) = d(y, x)$,
- 3) $d(x, z) \leq d(x, y) + d(y, z)$.

There is no condition that $d(x, y) = 0$ implies x and y coincide — this is where the adjective “pseudo” comes from, and the gadget is “extended” because we are allowing an infinite distance.

A metric space is an ep-metric space for which $d(x, y) = 0$ implies $x = y$, and all distances $d(x, y)$ are finite.

*Supported by NSERC.

The traditional objects of study in topological data analysis are finite metric spaces X , and the most common analysis starts by creating a family of simplicial complexes $V_s(X)$, the Vietoris-Rips complexes for X , which are parameterized by a distance variable s .

To construct the complex $V_s(X)$, it is harmless at the outset is to list the elements of X , or give X a total ordering — one can always do this without damaging the homotopy type. Then $V_s(X)$ is a simplicial complex (and a simplicial set), with simplices given by strings

$$x_0 \leq x_1 \leq \cdots \leq x_n$$

of elements of X such that $d(x_i, x_j) \leq s$ for all i, j . If $s \leq t$ then there is an inclusion $V_s X \subset V_t(X)$, and varying the distance parameter s gives a diagram (functor) $V_*(X) : [0, \infty] \rightarrow \mathbf{sSet}$, taking values in simplicial sets.

Following Spivak [4] (sort of), one can take an arbitrary diagram $Y : [0, \infty] \rightarrow \mathbf{sSet}$, and produce an ep-metric space $Re(Y)$, called its realization. This realization functor has a right adjoint S , called the singular functor, which takes an ep-metric space Z and produces a diagram $S(Z) : [0, \infty] \rightarrow \mathbf{sSet}$ in simplicial sets.

One needs good cocompleteness properties to construct the realization functor Re . Ordinary metric spaces are not well behaved in this regard, but it is shown in the first section (Lemma 3) that the category of ep-metric spaces has all of the colimits one could want. Then $Re(Y)$ can be constructed as a colimit of finite metric spaces U_s^n , one for each simplex $\Delta^n \rightarrow Y_s$ of some section of Y .

The metric space U_s^n is the set $\{0, 1, \dots, n\}$, equipped with a metric d , where $d(i, j) = s$ for $i \neq j$. A morphism $U_s^n \rightarrow Z$ of ep-metric spaces is a list $(x_0, x_1, \dots, x_n)$ of elements of Z such that $d(x_i, x_j) \leq s$ for all i, j . Such lists have nothing to with orderings on Z , and could have repeats.

With a bit of categorical homotopy theory, one shows (Proposition 7) that $Re(Y)$ is the set of vertices of the simplicial set Y_∞ (evaluation of Y at ∞), equipped with a metric that is imposed by the proof of Lemma 3.

One wants to know about the homotopy properties of the counit map $\eta : Y \rightarrow S(Re(Y))$, especially when Y is an old friend such as the Vietoris-Rips system $V_*(X)$. But $Re(V_*(X))$ is the original metric space X (Example 13), the object $S(X)$ is the diagram $[0, \infty] \rightarrow \mathbf{sSet}$ with $(S(X)_t)_n = \text{hom}(U_t^n, X)$, and the counit $\eta : V_t(X) \rightarrow S_t(X)$ in simplicial sets takes an n -simplex $\sigma : \Delta^n \rightarrow V_t(X)$ to the list $(\sigma(0), \sigma(1), \dots, \sigma(n))$ of its vertices.

We show in Section 3 (Proposition 16) that the map $\eta : V_t(X) \rightarrow S_t(X)$ is a weak equivalence for all distance parameter values t . The proof proceeds in two main steps, and involves technical results from the theory of simplicial approximation. The steps are the following:

1) We show (Lemma 14) that the map η induces a weak equivalence $\eta_* : BNV_t(X) \rightarrow BNS_t(X)$, where $\eta_* : NV_t(X) \rightarrow NS_t(X)$ is the induced comparison of posets of non-degenerate simplices. Here, $V_t(X)$ is a simplicial complex, so that $BNV_t(X)$ is a copy of the subdivision $\text{sd}(V_t(X))$, and is therefore weakly equivalent to $V_t(X)$.

2) There is a canonical map $\pi : \text{sd } S_t(X) \rightarrow BNS_t(X)$, and the second step in the proof of Proposition 16 is to show (Lemma 15) that this map π is a weak equivalence.

It follows that the map η induces a weak equivalence $\text{sd}(V_t(X)) \rightarrow \text{sd}(S_t(X))$, and Proposition 16 follows.

The fact that the space $S_t(X)$ is weakly equivalent to $V_t(X)$ for each t means that we have yet another diagram of spaces $S_*(X)$ that models persistent homotopy invariants for a data set X .

One should bear in mind, however, that $S_t(X)$ is an infinite complex. To see this, observe that if x_0 and x_1 are distinct points in X with $d(x_0, x_1) \leq t$, then all of the lists

$$(x_0, x_1, x_0, x_1, \dots, x_0, x_1)$$

define non-degenerate simplices of $S_t(X)$.

Contents

1	ep-metric spaces	3
2	Metric space realizations	7
2.1	The realization functor	8
2.2	Partial realizations	11
3	The singular functor	13

1 ep-metric spaces

An *extended pseudo-metric space* [2] (or an *uber metric space* [4]) is a set Y , together with a function $d : Y \times Y \rightarrow [0, \infty]$, such that the following conditions hold:

- a) $d(x, x) = 0$,
- b) $d(x, y) = d(y, x)$,
- c) $d(x, z) \leq d(x, y) + d(y, z)$.

Following [3], I use the term *ep-metric spaces* for these objects, which will be denoted by (Y, d) in cases where clarity is required for the metric.

Every metric space (X, d) is an ep-metric space, by composing the distance function $d : X \times X \rightarrow [0, \infty)$ with the inclusion $[0, \infty) \subset [0, \infty]$.

A morphism between ep-metric spaces (X, d_X) and (Y, d_Y) is a function $f : X \rightarrow Y$ such that

$$d_Y(f(x), f(y)) \leq d_X(x, y).$$

These morphisms are sometimes said to be non-expanding [2].

I shall use the notation $ep - \mathbf{Met}$ to denote the category of ep-metric spaces and their morphisms.

Example 1 (Quotient ep-metric spaces). Suppose that (X, d) is an ep-metric space and that $p : X \rightarrow Y$ is a surjective function.

For $x, y \in Y$, set

$$D(x, y) = \inf_P \sum_i d(x_i, y_i), \quad (1)$$

where each P consists of pairs of points (x_i, y_i) with $x = x_0$ and $y_k = y$, such that $p(y_i) = p(x_{i+1})$.

Certainly $D(x, x) = 0$ and $D(x, y) = D(y, x)$. One thinks of each P in the definition of $D(x, y)$ as a “polygonal path” from x to y . Polygonal paths concatenate, so that $D(x, z) \leq D(x, y) + D(y, z)$, and D gives the set Y an ep-metric space structure. This is the *quotient* ep-metric space structure on Y .

If x, y are elements of X , the pair (x, y) is a polygonal path from x to y , so that $D(p(x), p(y)) \leq d(x, y)$. It follows that the function p defines a morphism $p : (X, d) \rightarrow (Y, D)$ of ep-metric spaces.

Example 2 (Dividing by zero). Suppose that (X, d) is an ep-metric space. There is an equivalence relation on X , with $x \sim y$ if and only if $d(x, y) = 0$. Write $p : X \rightarrow X/\sim =: Y$ for the corresponding quotient map.

Given a polygonal path $P = \{(x_i, y_i)\}$ from x to y in X as above, $d(y_i, x_{i+1}) = 0$, so the sum corresponding to P in (1) can be rewritten as

$$d(x, y_0) + d(y_0, x_1) + d(x_1, y_1) + \cdots + d(x_k, y).$$

It follows that $d(x, y) \leq D(p(x), p(y))$, whereas $D(p(x), p(y)) \leq d(x, y)$ by construction.

Thus, if $D(p(x), p(y)) = 0$, then $d(x, y) = 0$ so that $p(x) = p(y)$.

Lemma 3. *The category $ep - \mathbf{Met}$ of ep-metric spaces is cocomplete.*

Proof. The empty set is the initial object for this category,

Suppose that $(X_i, d_i), i \in I$, is a list of ep-metric spaces. Form the set theoretic disjoint union $X = \sqcup_i X_i$, and define a function

$$d : X \times X \rightarrow [0, \infty]$$

by setting $d(x, y) = d_i(x, y)$ if x, y belong to the same summand X_i and $d(x, y) = \infty$ otherwise. Any collection of morphisms $f_i : X_i \rightarrow Y$ in $ep - \mathbf{Met}$ defines a unique function $f = (f_i) : X \rightarrow Y$, and this function is a morphism of $ep - \mathbf{Met}$ since

$$d(f(x), f(y)) = d(f_i(x), f_j(y)) \leq \infty = d(x, y)$$

if $x \in X_i$ and $y \in X_j$ with $i \neq j$.

Suppose given a pair of morphisms

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} X$$

in $ep - \mathbf{Met}$, and form the set theoretic coequalizer $\pi : X \rightarrow C$. The function p is the canonical map onto a set of equivalence classes of X , which classes are defined by the relations $f(a) \sim g(a)$ for $a \in A$. We give C the quotient ep-metric space structure, as in Example 1.

Suppose that $\alpha : (X, d_X) \rightarrow (Z, d_Z)$ is an morphism of ep-metric spaces such that $\alpha \cdot f = \alpha \cdot g$. Write $\alpha_* : C \rightarrow Z$ for the unique function such that $\alpha_* \cdot p = \alpha$.

Suppose given a polygonal path $P = \{(x_i, y_i)\}$ from x to y in X . Then $\alpha(y_i) = \alpha(x_{i+1})$, so that

$$d_Y(\alpha(x), \alpha(y)) \leq \sum_i d_Y(\alpha(x_i), \alpha(y_i)) \leq \sum_i d_X(x_i, y_i).$$

This is true for every polygonal path from x to y in X , so that

$$d_Y(\alpha_* p(x), \alpha_* p(y)) \leq d_C(p(x), p(y)).$$

It follows that $\alpha_* : (C, d_C) \rightarrow (Z, d_Z)$ is a morphism of ep-metric spaces. $\square$

Example 4 (“Bad” filtered colimit). If one starts with a diagram of metric spaces, the colimit C that is produced by Lemma 3 is an ep-metric space, and it may be that $d(x, y) = 0$ in the coequalizer C for some elements x, y with $x \neq y$.

In particular, suppose that $X_s = \{(\frac{1}{s\sqrt{2}}, 0), (0, \frac{1}{s\sqrt{2}})\} \subset \mathbb{R}^2$ for $0 < s < \infty$. Write $p_s = (\frac{1}{s\sqrt{2}}, 0)$ and $q_s = (0, \frac{1}{s\sqrt{2}})$ in X_s . Then $d(p_s, q_s) = \frac{1}{s}$. For $s \leq t$ there is an ep-metric space map $X_s \rightarrow X_t$ which is defined by $p_s \mapsto p_t$ and $q_s \mapsto q_t$.

The filtered colimit $\varinjlim_s X_s$ has two distinct points, namely p_∞ and q_∞ , and $d(p_\infty, q_\infty) \leq d(p_s, q_s) = \frac{1}{s}$ for all $s > 0$. It follows that $d(p_\infty, q_\infty) = 0$, whereas $p_\infty \neq q_\infty$.

Lemma 5. *Suppose that X is an ep-metric space. Then there is an isomorphism of ep-metric spaces*

$$\psi : \varinjlim_F F \xrightarrow{\cong} X,$$

where F varies over the finite subsets of X , with their induced ep-metric space structures.

Proof. The collection of finite subsets of X is filtered, and the set X is a filtered colimit of its finite subsets, so the function defining the ep-metric space map ψ is a bijection. Write d_∞ for the metric on the filtered colimit.

If $x, y \in X$ and $d(x, y) = s \leq \infty$ in X , then there is a finite subset F with $x, y \in F$ such that $d(x, y) = s$ in F . The list (x, y) is a polygonal path from x to y in F , so that $d_\infty(x, y) \leq d(x, y)$. It follows that $d(x, y) = d_\infty(x, y)$, and so ψ is an isomorphism. $\square$

An ep-metric space (X, d) has an associated system of posets $P_*(X) : [0, \infty] \rightarrow s\mathbf{Set}$, where $P_s(X)$ is the collection of finite subsets F of X such that $d(x, y) \leq s$ for any two members x, y of X .

This construction defines a system of abstract simplicial complexes $V_*(X)$, which can be constructed entirely within simplicial sets when X has a total ordering. In that case, the n -simplices of the simplicial set $V_s(X)$ are the strings $x_0 \leq x_1 \leq \dots \leq x_n$ such that $d(x_i, x_j) \leq s$. The diagram $V_*(X) : [0, \infty] \rightarrow s\mathbf{Set}$ is the Vietoris-Rips system. The spaces $V_s(X)$ are independent up to weak equivalence of the ordering on X , because there is a canonical weak equivalence (a “last vertex map”) $\gamma : BP_s(X) \rightarrow V_s(X)$ of systems, while the spaces $BP_s(X)$ are defined independently from the ordering. In classical terms, the nerve $BP_s(X)$ of the poset $P_s(X)$ (non-degenerate simplices of the Vietoris-Rips complex $V_s(X)$) is the barycentric subdivision of $V_s(X)$.

Example 6 (Excision for path components). Suppose that X and Y are finite subsets of an ep-metric space Z , with the induced ep-metric space structures. Consider the inclusions of finite ep-metric spaces

$$\begin{array}{ccc} X \cap Y & \longrightarrow & Y \\ \downarrow & & \downarrow \\ X & \longrightarrow & X \cup Y \end{array}$$

inside Z . Write $X \cup_m Y$ for the corresponding pushout in the category of ep-metric spaces. The unique map

$$X \cup_m Y \rightarrow X \cup Y$$

of ep-metric spaces is the identity on the underlying point set. Write d_m for the metric on $X \cup_m Y$. Then $d_m(x, y)$ is the minimum of sums

$$\sum d(x_i, x_{i+1}), \tag{2}$$

indexed over paths

$$P : x = x_0, x_1, \dots, x_n = y,$$

such that for each i the points x_i, x_{i+1} are either both in X or both in Y .

All sums in (2) are finite, and $d_m(x, y)$ is realized by a particular path P since X and Y are finite. Note that $d(x, y) \leq d_m(x, y)$, by construction, and that $d(x, y) = d_m(x, y)$ if x, y are both in either X or Y .

There are induced simplicial set maps

$$V_s(X) \cup V_s(Y) \rightarrow V_s(X \cup_m Y) \rightarrow V_s(X \cup Y),$$

all of which are the identity on vertices. There is a 1-simplex $\sigma = \{x, y\}$ of $V_s(X \cup_m Y)$ if and only if there is a path

$$P : x = x_0, x_1, \dots, x_n = y$$

consisting of 1-simplices in either X or Y , such that

$$\sum d(x_i, x_{i+1}) \leq s.$$

Then all $d(x_i, x_{i+1}) \leq s$, so that x and y are in the same path component of $V_s(X) \cup V_s(Y)$. It follows that there is an induced isomorphism

$$\pi_0(V_s(X) \cup V_s(Y)) \cong \pi_0 V_s(X \cup_m Y). \quad (3)$$

The isomorphisms (3) induce isomorphisms

$$\pi_0(V_s(X) \cup V_s(Y)) \cong \pi_0 V_s(X \cup_m Y). \quad (4)$$

for arbitrary subsets X and Y of an ep-metric space Z , by an application of Lemma 5.

2 Metric space realizations

Write U_s^n for the collection of axis points $x_i = \frac{s}{\sqrt{2}}e_i$, where

$$e_i = (0, \dots, \overset{i+1}{1}, \dots, 0) \in \mathbb{R}^{n+1}.$$

for $0 \leq i \leq n$. Observe that $d(x_i, x_j) = s$ in $\mathbb{R}^{n+1}$ for $i \neq j$. Another way of looking at it: U_s^n is the set $\mathbf{n} = \{0, 1, \dots, n\}$ with $d(i, j) = s$ for $i \neq j$.

An ep-metric space morphism $f : U_s^n \rightarrow Y$ consists of points $f(x_i)$, $0 \leq i \leq n$, such that $d_Y(f(x_i), f(x_j)) \leq s$ for all i, j .

Write $s\mathbf{Set}^{[0, \infty]}$ for the category of diagrams (functors) $X : [0, \infty] \rightarrow s\mathbf{Set}$ and their natural transformation, which take values in simplicial sets and are defined on the poset $[0, \infty]$. I usually write $s \mapsto X_s$ for such a diagram X . In particular, X_∞ is the value that the diagram X takes at the terminal object of $[0, \infty]$.

Suppose that K is a simplicial set. The representable diagram $L_s K$ satisfies the universal property

$$\mathrm{hom}(L_s K, X) \cong \mathrm{hom}(K, X_s).$$

One shows that

$$(L_s K)_t = \begin{cases} \emptyset & \text{if } t < s, \\ K & \text{if } t \geq s. \end{cases}$$

The set of maps $L_s \Delta^n \rightarrow X$ can be identified with the set of n -simplices of the simplicial set X_s .

A morphism $L_t \Delta^m \rightarrow L_s \Delta^n$ consists of a relation $s \leq t$ and a simplicial map $\theta : \Delta^m \rightarrow \Delta^n$. In the presence of such a morphism, the function $\theta : \mathbf{m} \rightarrow \mathbf{n}$ defines an ep-metric space morphism $U_t^m \rightarrow U_s^n$, since $s = d(\theta(i), \theta(j)) \leq d(i, j) = t$.

2.1 The realization functor

Suppose that $X : [0, \infty] \rightarrow s\mathbf{Set}$ is a diagram. The category $\mathbf{\Delta}/X$ of simplices of X has maps $L_s \Delta^n \rightarrow X$ as objects and commutative diagrams

$$\begin{array}{ccc} L_t \Delta^m & \xrightarrow{\theta} & L_s \Delta^n \\ & \searrow \tau & \swarrow \sigma \\ & X & \end{array}$$

as morphisms.

Equivalently, a simplex of X is a simplicial set map $\Delta^n \rightarrow X_s$, and a morphism of simplices is a diagram

$$\begin{array}{ccc} \Delta^m & \xrightarrow{\theta} & \Delta^n \\ \tau \downarrow & & \downarrow \sigma \\ X_t & \longleftarrow & X_s \end{array} \quad (5)$$

Every simplex $\Delta^n \rightarrow X_s$ determines a simplex

$$\Delta^n \rightarrow X_s \rightarrow X_\infty,$$

and we have a functor $r : \mathbf{\Delta}/X \rightarrow \mathbf{\Delta}/X_\infty$, where $\mathbf{\Delta}/X_\infty$ is the simplex category of the simplicial set X_∞ .

There is an inclusion $i : \mathbf{\Delta}/X_\infty \rightarrow \mathbf{\Delta}/X$, and the composite $r \cdot i$ is the identity. The maps

$$\begin{array}{ccc} \Delta^n & \xrightarrow{1} & \Delta^n \\ r(\sigma) \downarrow & & \downarrow \sigma \\ X_\infty & \longleftarrow & X_s \end{array} \quad (6)$$

define a natural transformation $h : i \cdot r \rightarrow 1$.

There is a functor

$$\mathbf{\Delta}/X \rightarrow s\mathbf{Set} \quad (7)$$

which takes a morphism (5) to the map $\theta : \Delta^m \rightarrow \Delta^n$.

The translation category E_X for the functor (7) is a simplicial category that has objects consisting of pairs (σ, x) where $\sigma : \Delta^n \rightarrow X_s$ and $x \in \Delta^n$ (of a fixed dimension). A morphism $(\tau, y) \rightarrow (\sigma, x)$ of E_X is a morphism $\theta : \tau \rightarrow \sigma$ as in (5) such that $\theta(y) = x$. The path component simplicial set $\pi_0 E_X$ of the category E_X is isomorphic to the colimit

$$\varinjlim_{L_s \Delta^n \rightarrow X} \Delta^n.$$

There is a corresponding translation category E_{X_∞} for the functor which takes the simplex $\Delta^n \rightarrow X_\infty$ to the simplicial set Δ^n , and there is an induced

functor $i_* : E_{X_\infty} \subset E_X$. The functor $r : \mathbf{\Delta}/X \rightarrow \mathbf{\Delta}/X_\infty$ induces a functor $r_* : E_X \rightarrow E_{X_\infty}$. The composite $r_* \cdot i_*$ is the identity on E_{X_∞} , and the map (6) defines a natural transformation $i_* \cdot r_* \rightarrow 1$ of functors $E_{X_\infty} \rightarrow E_{X_\infty}$.

The translation categories E_X and E_{X_∞} are therefore homotopy equivalent, and thus have isomorphic simplicial sets of path components. It follows that there are isomorphisms of simplicial sets

$$X_\infty \xleftarrow{\cong} \varinjlim_{\Delta^n \rightarrow X_\infty} \Delta^n \xrightarrow{\cong} \varinjlim_{L_s \Delta^n \rightarrow X} \Delta^n \quad (8)$$

Suppose that $X : [0, \infty] \rightarrow \mathbf{sSet}$ is a diagram, and set

$$Re(X) = \varinjlim_{L_s \Delta^n \rightarrow X} U_s^n$$

in the category of ep-metric spaces.

It follows from the identifications of (8) that $Re(X)$ is the set of vertices of X_∞ , equipped with an ep-metric space structure.

If x and y are two such vertices, and are the boundary of a 1-simplex

$$\Delta^1 \rightarrow X_s \rightarrow X_\infty$$

then x and y are in the image of a map $U_s^1 \rightarrow Re(X)$, so that $d(x, y) \leq s$. If there is a sequence of 1-simplices $\omega_i : \Delta^1 \rightarrow X_{s_i}$ that define a polygonal path

$$P : x = x_0 \hookrightarrow x_1 \hookrightarrow \cdots \hookrightarrow x_k = y$$

of 1-simplices in X_∞ , then $d(x, y) \leq \sum_i s_i$ by definition. Formally, we set

$$d(x, y) = \inf_P \left\{ \sum_i s_i \right\}. \quad (9)$$

provided such polygonal paths exist. Otherwise, we set $d(x, y) = \infty$.

The resulting metric d is the metric which is imposed on the set of vertices of X_∞ by the requirement that

$$Re(X) = \varinjlim_{L_s \Delta^n \rightarrow X} U_s^n$$

in the category of ep-metric spaces — see Lemma 3. We have shown the following:

Proposition 7. *Suppose that $X : [0, \infty] \rightarrow \mathbf{sSet}$ is a diagram. Then the ep-metric space $Re(X)$ has underlying set given by the set of vertices of X_∞ , with metric defined within path components by (9). Elements x and y that are in distinct path components have $d(x, y) = \infty$.*

Example 8 (Realization of Vietoris-Rips systems). Suppose that X is a finite ep-metric space, and that X is totally ordered.

The realization $Re(V_*(X))$ has X as its underlying set, and $V_\infty(X) = \Delta^X$ is a finite simplex, which is connected, so that there is a finite polygonal path in X between any two points $x, y \in X$. We have a relation

$$d(x, y) \leq \sum_i s_i$$

in X for any polygonal path P which is defined by 1-simplices $\omega_i \in V_{s_i}(X)$. This means that $d(x, y)$ in X coincides with the distance between x and y in the metric space $Re(X)$. It follows that the identity on the set X induces an isomorphism of ep-metric spaces

$$\phi : Re(V_*(X)) \xrightarrow{\cong} X.$$

This map ϕ is an isomorphism of ep-metric spaces, by Lemma 5 and the previous paragraphs.

Example 9 (Degree Rips systems). Continue with a finite totally ordered ep-metric space X as in Example 8, let k be a positive integer, and consider the degree Rips system $L_{*,k}(X)$. We choose k such that the system of complexes $L_{*,k}(X)$ is non-empty, ie. such that $k \leq |X|$. Then $L_{t,k}(X) = V_t(X)$ for t sufficiently large, and X is the underlying set of $Re(L_{*,k}(X))$.

The maps

$$Re(L_{*,k}(X)) \rightarrow Re(V_*(X)) \rightarrow X$$

are isomorphisms of metric spaces, by a cofinality argument.

Here is a special case:

Lemma 10. *Suppose that K is a simplicial complex and that $s > 0$. Then $Re(L_s K) = Re(L_s \text{sk}_1(K))$, and $Re(L_s K)$ is the set of vertices K_0 with a metric d defined by*

$$d(x, y) = \begin{cases} \infty & \text{if } [x] \neq [y] \text{ in } \pi_0(K), \\ \min_P s \cdot k & \text{if } [x] = [y]. \end{cases}$$

where P varies through the polygonal paths

$$P : x = x_0 \leftrightsquigarrow x_1 \leftrightsquigarrow \cdots \leftrightsquigarrow x_k = y$$

of 1-simplices between x and y .

Proof. Write $Re(K) = Re(L_s K)$. The simplicial set K is a colimit of its simplices, and so there is an isomorphism

$$\varinjlim_{\Delta^n \rightarrow K} L_s \Delta^n \xrightarrow{\cong} L_s K.$$

It follows that there is an isomorphism

$$\varinjlim_{\Delta^n \rightarrow K} U_s^n \xrightarrow{\cong} Re(L_s K).$$

Suppose that $n \geq 2$. Then $\partial\Delta^n$ and Δ^n have the same vertices, and any two vertices x, y are on a common face $\Delta^{n-1} \subset \Delta^n$. It follows that $d(x, y) = s$ in $Re(\partial\Delta^n)$ and $Re(\Delta^n)$, and the induced map

$$Re(\partial\Delta^n) \rightarrow Re(\Delta^n)$$

is an isomorphism for $n \geq 2$.

The displayed metric d on the vertices of K defines a metric space $Re(K)$, with maps $\sigma_* : U_s^n \rightarrow Re(K)$ for all simplices $\sigma : \Delta^n \rightarrow K$, which maps are natural with respect to the simplicial structure of K .

Any family of metric space morphisms $f_\sigma : U_s^n \rightarrow Y$ determines a unique function $f : K_0 \rightarrow Y$. Also, $d(f(x), f(y)) \leq s$ if x, y are in a common simplex $\Delta^1 \rightarrow K$. If P is a polygonal path between x and y as above, then $d(f(x), f(y)) \leq k \cdot s$. This is true for all such polygonal paths, so $d(f(x), f(y)) \leq d(x, y)$.

If x and y are in distinct components of K , then $d(f(x), f(y)) \leq d(x, y) = \infty$. $\square$

The following result says that the realization $Re(X)$ of a diagram $X : [0, \infty] \rightarrow s\mathbf{Set}$ depends only on the associated diagram of graphs $sk_1(X)$.

Lemma 11. *Suppose that $X : [0, \infty] \rightarrow s\mathbf{Set}$ is a diagram. Then the inclusion $sk_1 X \subset X$ induces an isomorphism*

$$Re(sk_1 X) \xrightarrow{\cong} Re(X).$$

Proof. The diagram of 1-skeleta $sk_1 X$ is a colimit

$$\varinjlim_{L_s \Delta^n \rightarrow X} L_s sk_1 \Delta^n,$$

since the functor sk_1 preserves colimits. There are commutative diagrams

$$\begin{array}{ccc} Re(L_s sk_1 \Delta^n) & \longrightarrow & Re(sk_1 X) \\ \cong \downarrow & & \downarrow \\ Re(L_s \Delta^n) & \longrightarrow & Re(X) \end{array}$$

that are natural in the simplices of X , and it follows that the induced map $Re(sk_1 X) \rightarrow Re(X)$ is an isomorphism, as required. $\square$

2.2 Partial realizations

Suppose again that $X : [0, \infty] \rightarrow s\mathbf{Set}$ is a diagram in simplicial sets. We construct partial realizations by writing

$$Re(X)_s = \varinjlim_{L_t \Delta^n \rightarrow X, t \leq s} U_t^n.$$

This is the colimit of a functor taking values in ep-metric spaces, which is defined on the full subcategory $\Delta/X_{\leq s}$ of Δ/X having objects $L_t\Delta^n \rightarrow X$ with $t \leq s$. A map $L_t\Delta^n \rightarrow X$ can be identified with a simplex $\Delta^n \rightarrow X_t$, and the relation $t \leq s$ defines a simplex $\Delta^n \rightarrow X_t \rightarrow X_s$, so that we have a functor

$$r : \Delta/X_{\leq s} \rightarrow \Delta/X_s,$$

along with an inclusion $i : \Delta/X_s \subset \Delta/X_{\leq s}$. The composite $r \cdot i$ is the identity, and the composite $i \cdot r$ is homotopic to the identity, just as before.

There is a functor $\Delta/X_{\leq s} \rightarrow s\mathbf{Set}$ which takes a simplex $\Delta^n \rightarrow X_t$ to the simplicial set Δ^n . By manipulating path components of homotopy colimits, one finds isomorphisms

$$X_s \xleftarrow{\cong} \varinjlim_{\Delta^n \rightarrow X_s} \Delta^n \xrightarrow{\cong} \varinjlim_{L_t\Delta^n \rightarrow X, t \leq s} \Delta^n.$$

that are analogous to the isomorphisms of (9). It follows, as in Proposition 7, that the set underlying the metric space $Re(X)_s$ is the set of vertices of the simplicial set X_s .

The metric d on $(X_s)_0$ is defined as before: $d(x, y) = \infty$ if x and y not in the same path component of X_s . Otherwise

$$d(x, y) = \inf_P \left\{ \sum t_i \right\}, \quad (10)$$

indexed over all polygonal paths

$$P : x = x_0 \rightrightarrows x_1 \rightrightarrows \cdots \rightrightarrows x_k = y$$

that are defined by 1-simplices $\omega : \Delta^1 \rightarrow X_{t_i}$ with $t_i \leq s$.

We then have the following analogue of Proposition 7:

Proposition 12. *Suppose that $X : [0, \infty] \rightarrow s\mathbf{Set}$ is a functor. Then the ep-metric space $Re(X)_s$ has underlying set given by the set of vertices of X_s , with metric defined within path components by (10). Elements x, y in distinct path components have $d(x, y) = \infty$.*

The map $X_s \rightarrow X_\infty$ defines a map $Re(X)_s \rightarrow Re(X)$ and $Re(X)_\infty = Re(X)$. There is an isomorphism of ep-metric spaces

$$\varinjlim_s Re(X)_s \xrightarrow{\cong} Re(X), \quad (11)$$

since the element ∞ is terminal in $[0, \infty]$ and $Re(X)_\infty = Re(X)$.

Example 13 (Partial metrics for Vietoris-Rips complexes). Suppose that X is a finite totally ordered ep-metric space. Consider the associated functor $V_*(X) : [0, \infty] \rightarrow s\mathbf{Set}$.

The associated ep-metric space $Re(X)_s$ has underlying set X . We have $d(x, y) = \infty$ if x, y are in distinct path components of $V_s(X)$. Otherwise

$$d(x, y) = \inf_P \left\{ \sum d(x_i, x_{i+1}) \right\},$$

indexed over all polygonal paths

$$P : x = x_0, x_1, \dots, x_n = y,$$

with $d(x_i, x_{i+1}) \leq s$.

If $d(x, y) = t \leq s$ in X then $d(x, y) = t$ in $Re(X)_s$. Otherwise, the distance between x and y in the same path component of $Re(X)_s$ is more interesting — it is achieved by a particular path P since X is finite, and $d(x, y)$ is a type of weighted path length.

We see in Example 8 that there is an isomorphism of ep-metric spaces $\phi : Re(V_*(X)) \xrightarrow{\cong} X$. It follows that there is an ep-metric space map $\phi_s : Re(X)_s \rightarrow X$ which is the identity on the underlying point set X , and compresses distances.

3 The singular functor

The right adjoint S of the realization functor Re is defined for an ep-metric space Y by

$$S(Y)_{s,n} = \text{hom}(U_s^n, Y),$$

where $\text{hom}(U_s^n, Y)$ is the collection of ep-metric space morphisms $U_s^n \rightarrow Y$. Equivalently, $S(Y)_{s,n}$ is the set of families of points $(x_0, x_1, \dots, x_n)$ in Y such that $d(x_i, x_j) \leq s$.

A simplex $(x_0, x_1, \dots, x_n)$ is alternatively a function $\mathbf{n} \rightarrow Y$ (a “bag of words”), with a distance restriction. There is no requirement that the elements x_i are distinct. This simplex is non-degenerate if and only if $x_i \neq x_{i+1}$ for $0 \leq i \leq n-1$.

Suppose that an ep-metric space X is totally ordered, as in Example 8 above. Then, in view of the discussion of Example 8, the canonical map $\eta : V_*(X) \rightarrow SRe(V_*(X))$ consists of functions $\eta : V_t(X) \rightarrow S_t(X)$ which send simplices $\sigma : x_0 \leq x_1 \leq \dots \leq x_n$ with $d(x_i, x_j) \leq t$ to the list of points $(x_0, x_1, \dots, x_n)$.

If σ is non-degenerate, so that the vertices x_i are distinct, then $\eta(\sigma)$ is a non-degenerate simplex of $S_t(X)$.

The poset NZ of non-degenerate simplices of a simplicial set Z has $\sigma \leq \tau$ if there is a subcomplex inclusion $\langle \sigma \rangle \subset \langle \tau \rangle$, where $\langle \sigma \rangle$ is the subcomplex of Z which is generated by the simplex σ . Equivalently, $\sigma \leq \tau$ if there is an ordinal number map θ such that $\theta^*(\tau) = \sigma$.

The map η induces a morphism $\eta_* : NV_t(X) \rightarrow NS_t(X)$ of posets of non-degenerate simplices.

Lemma 14. *Suppose that X is a totally ordered ep-metric space. Then the induced simplicial set map*

$$\eta_* : BNV_t(X) \rightarrow BNS_t(X)$$

of associated nerves is a weak equivalence.

Proof. Given a non-degenerate simplex $\sigma \in S_t(X)$, write $L(\sigma)$ for its list of distinct elements.

Suppose that $\langle \tau \rangle \subset \langle \sigma \rangle$, where τ and σ are non-degenerate simplices of $S_t(X)$. Then $\tau = s \cdot d(\sigma)$ for an (iterated) face map d and degeneracy s . Then

$$L(\tau) = L(s \cdot d(\sigma)) = L(d(\sigma)) \subset L(\sigma).$$

It follows that the assignment $\sigma \mapsto L(\sigma)$ defines a poset morphism

$$L : NS_t(X) \rightarrow NV_t(X).$$

The composite

$$NV_t(X) \xrightarrow{\eta} NS_t(X) \xrightarrow{L} NV_t(X)$$

is the identity on $NV_t(X)$.

Consider the composite poset morphism

$$NS_t(X) \xrightarrow{L} NV_t(X) \xrightarrow{\eta} NS_t(X). \quad (12)$$

Given a non-degenerate simplex $\tau = (y_0, \dots, y_r)$ of $S_t(X)$, write $L(\tau) = (s_0, \dots, s_k)$ for the list of distinct elements of τ , in the order specified by the total order for X . Then the list

$$V(\tau) = (y_0, \dots, y_r, s_0, \dots, s_k)$$

is a simplex of $S_t(X)$, since each s_j is some y_{i_j} , and there are relations

$$\langle \tau \rangle \leq \langle V(\tau) \rangle \geq \langle L(\tau) \rangle$$

as subcomplexes of $S_t(X)$.

The simplex $V(\tau)$ has the form $V(\tau) = s(V_*(\tau))$ for a unique iterated degeneracy s and a unique non-degenerate simplex $V_*(\tau)$ (see Lemma 18), and $\langle V(\tau) \rangle = \langle V_*(\tau) \rangle$.

Suppose that γ is non-degenerate in $S_t(X)$ and that $\gamma \in \langle \tau \rangle$. Then $\gamma = d(\tau)$ for some face map d , and $\gamma = (x_0, \dots, x_k)$ is a sublist of $\tau = (y_0, \dots, y_r)$. The ordered list $L(\gamma)$ of distinct elements of γ is a sublist of $L(\tau)$, and $V(\gamma)$ is a sublist of $V(\tau)$. There is a diagram of relations

$$\begin{array}{ccccc} \langle \tau \rangle & \longrightarrow & \langle V_*(\tau) \rangle & \longleftarrow & \langle L(\tau) \rangle \\ \uparrow & & \uparrow & & \uparrow \\ \langle \gamma \rangle & \longrightarrow & \langle V_*(\gamma) \rangle & \longleftarrow & \langle L(\gamma) \rangle \end{array}$$

It follows that the composite (12) is homotopic to the identity on the poset $NS_t(X)$, and the Lemma follows. $\square$

The subdivision $\text{sd}(Z)$ of a simplicial set Z is defined by

$$\text{sd}(Z) = \varinjlim_{\Delta^n \rightarrow Z} BN\Delta^n.$$

The poset morphisms $N\Delta^n \rightarrow NZ$ that are induced by simplices $\Delta^n \rightarrow Z$ together induce a map

$$\pi : \text{sd}(Z) \rightarrow BNZ.$$

It is known [1] (and not difficult to prove) that the map π is a bijection for simplicial sets Z that are polyhedral.

A polyhedral simplicial set is a subobject of the nerve of a poset. All oriented simplicial complexes are polyhedral in this sense. Examples include the Vietoris-Rips systems $V_s(X)$ associated to a totally ordered ep-metric space X , since

$$V_s(X) \subset V_\infty(X) = BX,$$

where BX is the nerve of the totally ordered poset X .

Lemma 15. *Suppose that X is an ep-metric space. Then the map*

$$\pi : \text{sd}(S_t(X)) \rightarrow BNS_t(X)$$

is a weak equivalence.

Proof. We show that all subcomplexes $\langle \sigma \rangle$ which are generated by non-degenerate simplices σ of $S_t(X)$ are contractible. Then Lemma 4.2 of [1] implies that the map π is a weak equivalence.

A non-degenerate simplex σ has the form $\sigma = (x_0, x_1, \dots, x_k)$ with $x_i \neq x_{i+1}$. The simplices τ of $\langle \sigma \rangle$ have the form

$$\tau = \theta^* \sigma = (x_{\theta(0)}, \dots, x_{\theta(k)}),$$

where $\theta : \mathbf{k} \rightarrow \mathbf{n}$ is an ordinal number morphism.

For each such θ , the list

$$(x_0, x_{\theta(0)}, \dots, x_{\theta(k)})$$

defines a simplex τ_* of $\langle \sigma \rangle$, since τ_* is a face of the simplex

$$s_0(\sigma) = (x_0, x_0, \dots, x_k).$$

The simplices τ_* define functors

$$\begin{array}{ccccccc} x_0 & \longrightarrow & x_0 & \longrightarrow & \dots & \longrightarrow & x_0 \\ \downarrow & & \downarrow & & & & \downarrow \\ x_{\theta(0)} & \longrightarrow & x_{\theta(1)} & \longrightarrow & \dots & \longrightarrow & x_{\theta(m)} \end{array}$$

or homotopies, that consist of simplices of $\langle \sigma \rangle$ that patch together to give a contracting homotopy $\langle \sigma \rangle \times \Delta^1 \rightarrow \langle \sigma \rangle$. $\square$

Proposition 16. *Suppose that X is a totally ordered ep-metric space. Then there is a diagram of weak equivalences*

$$\begin{array}{ccccc} BNV_t(X) & \xleftarrow[\cong]{\pi} & \text{sd}(V_t(X)) & \xrightarrow{\gamma} & V_t(X) \\ \eta_* \downarrow & & \downarrow \eta_* & & \downarrow \eta \\ BNS_t(X) & \xleftarrow[\pi]{} & \text{sd}(S_t(X)) & \xrightarrow[\gamma]{} & S_t(X) \end{array}$$

In particular, the map $\eta : V_t(X) \rightarrow S_t(X)$ is a weak equivalence.

This diagram is natural in t .

Proof. The map $\eta_* : BNV_t(X) \rightarrow BNS_t(X)$ is a weak equivalence by Lemma 14. The map $\pi : \text{sd}(S_t(X)) \rightarrow BNS_t(X)$ is a weak equivalence by Lemma 15. The instances of the maps γ are weak equivalences [1]. It follows that the maps $\eta_* : \text{sd}(V_t(X)) \rightarrow \text{sd}(S_t(X))$ and $\eta : V_t(X) \rightarrow S_t(X)$ are weak equivalences. $\square$

Remark 17. The total ordering on the ep-metric space X in Proposition 16 is intimately involved in the definition of the Vietoris-Rips system $V_*(X)$, the morphism

$$\eta : V_t(X) \rightarrow S_t(\text{Re}(V_*(X))) = S_t(X),$$

and all induced maps η_* .

The counit $\eta : Z \rightarrow S(\text{Re}(Z))$ is not a sectionwise weak equivalence in general. One can show that $S_\infty(\text{Re}(Z))$ is the nerve of the trivial groupoid on the vertex set of Z_∞ (Proposition 7), and is therefore contractible, whereas the space Z_∞ may not be contractible. For example, if K is a simplicial set, then there is an identification $K = (L_s K)_\infty$.

It may be that the map

$$\eta : BP_t(X) \rightarrow S_t(\text{Re}(BP_*(X)))$$

is a weak equivalence for arbitrary ep-metric spaces X , but this has not been proved. Such a result would give a non-oriented version of Proposition 16.

The following is a classical result, which is included here for the sake of completeness. This result is usually neither expressed nor proved in the form displayed here.

Lemma 18. *Suppose that σ is an n -simplex of a simplicial set X . Then there is a unique iterated degeneracy and a non-degenerate simplex x such that $\sigma = s(x)$.*

An iterated degeneracy is a surjective ordinal number map $s : \mathbf{n} \rightarrow \mathbf{k}$. Such a map induces a function $s : X_k \rightarrow X_n$ for a simplicial set X . Lemma 18 says that $\sigma = s(x)$ for some iterated degeneracy s and a non-degenerate simplex x , and that this representation is unique.

Proof of Lemma 18. Suppose that $\sigma = s(x) = s'(x')$ where s, s' are iterated degeneracies and x, x' are non-degenerate.

The $x = d(\sigma)$ for some face map d such that $d \cdot s = 1$, and so $d(s'(x')) = s''(d''(x))$ for some iterated degeneracy s'' and face map d'' . But x is non-degenerate, so that $s'' = 1$ and $x = d''(x')$. Similarly, $x' = \tilde{d}(x)$ for some face map $\tilde{d}$. But then x and x' have the same dimension, and $d = 1$, so that $x = x'$.

If $s \neq s'$ there is a face map d such that $d \cdot s = 1$ but $d \cdot s' \neq 1$. Then $\sigma = s(x) = s'(x)$ for $s \neq s'$ and x non-degenerate, then

$$d(\sigma) = x = d(s'(x)) = s''(d''(x))$$

for some degeneracy s'' and face map d'' , at least one of which is non-trivial.

But x is non-degenerate, so that $s'' = 1$ and $x = d''(x)$ only if $d'' = 1$. This contradicts the assumption that $s \neq s'$. $\square$

References

- [1] J. F. Jardine. Simplicial approximation. *Theory Appl. Categ.*, 12:No. 2, 34–72 (electronic), 2004.
- [2] Leland McInnes and John Healy. UMAP: uniform manifold approximation and projection for dimension reduction. *CoRR*, abs/1802.03426, 2018.
- [3] Luis N. Scoccola. *Locally Persistent Categories and Metric Properties of Interleaving Distances*. Scholarship@Western, 2020. Thesis (Ph.D.), University of Western Ontario.
- [4] D.I. Spivak. Metric realization of fuzzy simplicial sets. Preprint, 2009.